



Tavitian-Exley, I., Maheu-Giroux, M., Platt, L., Heimer, R., Uusküla, A., Levina, O., ... Boily, M. C. (2018). Differences in risk behaviours and HIV status between primary amphetamines and opioid injectors in Estonia and Russia. *International Journal of Drug Policy*, 53, 96-105.
<https://doi.org/10.1016/j.drugpo.2017.11.010>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.drugpo.2017.11.010](https://doi.org/10.1016/j.drugpo.2017.11.010)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <https://www.sciencedirect.com/science/article/pii/S0955395917303390> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms>

Differences in risk behaviours and HIV status between primary amphetamines and opioid injectors in Estonia and Russia

Authors: Isabel TAVITIAN-EXLEY^{1*}; Mathieu MAHEU-GIROUX²; Lucy PLATT³; Robert HEIMER⁴; Anneli UUSKÜLA⁵; Olga LEVINA⁶; Peter VICKERMAN⁷; Marie-Claude BOILY^{1*}

¹ Department of Infectious Disease Epidemiology, Imperial College London, United Kingdom.

² Department of Epidemiology, Biostatistics, and Occupational Health, McGill University, Montréal, Canada.

³ Department of Social and Environmental Health Research, London School of Hygiene and Tropical Medicine.

⁴ Epidemiology of Microbial Diseases, School of Public Health, Yale University, United States.

⁵ Department of Family Medicine and Public Health, University of Tartu, Tartu, Estonia.

⁶ NGO Stellit, St Petersburg, Russian Federation

⁷ School of Social and Community Medicine, University of Bristol, Bristol, United Kingdom.

*Corresponding Author:

Tavitianexley@gmail.com (i.tavitianexley09@imperial.ac.uk) and mc.boily@imperial.ac.uk

Submission category: Original research article

Running head: Amphetamine injection and HIV risk

Word count: 4,100 (excluding title page, references, figures and tables)

Keywords: HIV; risk behaviours; PWID; amphetamine injection; opioid injection; polydrug use; respondent-driven sampling; hepatitis C; herpes simplex virus.

Declaration of interest: None declared.

ABSTRACT

Background and objective

People who inject drugs (PWID) account for over half of new HIV infections in Eastern Europe and central Asia, where opioids continue to be the dominant illicit drugs injected. Stimulants including amphetamines (ATS) have been associated with HIV infection risk in several settings. We sought to examine whether primary ATS injection was associated with greater HIV risk, compared to opioid injection in two European locales with significant HIV epidemics.

Methods

PWID in Kohtla-Järve and St. Petersburg were recruited using respondent-driven sampling in 2012-2013. Survey data on demographic characteristics, service use, injecting and sexual risk behaviours; and HIV-status (and HCV in Kohtla-Järve) were compared between primary opioid and ATS injectors using logistic regression models.

Results

Of 591 injectors recruited in Kohtla-Järve and 811 in St. Petersburg, 195 (33%) and 27 (4%) primarily injected ATS in each city. In both cities, ATS injectors were younger than opioid injectors, initiated injection later, injected less frequently and were more likely to have been paid for sex. In both cities, PWID had high levels of multiple sex partners. In Kohtla-Järve, ATS-injectors had lower odds of back-loading and greater odds of polydrug use than opioid-injectors. In St. Petersburg, where over half of PWID reported unsafe sharing practices, ATS-injectors were less likely to report these practices. ATS-injection was negatively associated with being HIV positive in Kohtla-Järve (aOR=0.6; 95%CI: 0.5-0.8) and St. Petersburg (aOR=0.3; 95%CI: 0.1-0.7). ATS-injection was negatively associated with HCV-reactivity in Kohtla-Järve (aOR=0.5; 95%CI: 0.3-0.6).

Conclusions

In both locations, primary ATS injection was associated with lower injecting risk behaviours, lower odds of HIV and being paid for sex compared to opioid injection. Interventions targeting the characteristics and needs of ATS injectors are needed to increase contact with services and reduce sexual and injecting risk. Harm reduction services, including sexual risk reduction, need to be expanded for all PWID in St. Petersburg.

INTRODUCTION

People who inject drugs (PWID) were estimated to account for 51% of new human immunodeficiency virus (HIV-1) infections in Eastern Europe and central Asia in 2014, a region with the fastest growing HIV epidemic associated with injection drug use globally (UNAIDS, 2016; UNODC, 2016).

Opioid injection has been the main driver of HIV epidemics in Estonia and the Russian Federation, where over half of PWID in Kohtla-Järve (Estonia) and St. Petersburg (Russian Federation) were seropositive in 2012 (El-Bassel, et al., 2013; Jolley, et al., 2012; Uusküla, Raag, et al., 2015; Walsh, et al., 2013). Both cities are situated on the Baltic Sea, on the northern part of two major heroin trafficking corridors linking Afghanistan to the heroin markets of Western Europe; both have experienced HIV epidemics driven by transmission among PWID since the late 1990s (UNODC, 2012, 2015b). Despite similar HIV prevalence, estimated incidence was higher in Kohtla-Järve (22/100 person-years, 2012) than in St. Petersburg (7.2/100 person-years, 2010), partly due to the higher proportion of young PWID in the Estonian city and to differences in the context and epidemic response, presented in Table 1. Evidence-based harm reduction interventions, including needle and syringe programmes (NSP) and opiate substitution treatment (OST), were introduced in Kohtla-Järve in 2004 (Estonia Ministry of Health, 2014; Mathers, et al., 2010) whereas in St Petersburg, OST remains illegal and clean needles and syringes are not endorsed by government and provided by a few non-governmental organisations (Degenhardt, et al., 2014; EMCDDA, 2015).

Most PWID injected heroin and illicitly-produced synthetic opioids, namely fentanyl in Kohtla-Järve (introduced into Estonia following a heroin shortage in 2000) and methadone in St Petersburg (Eritsyan, et al., 2013; Heimer, et al., 2015). Estonia has reported one of the highest prevalences of amphetamine type stimulant (ATS) use in Europe (EMCDDA, et al., 2012; UNODC, 2014) and ATS have emerged as a major secondary drug among PWID in Kohtla-Järve and St. Petersburg (EMCDDA, 2010; Grund, et al., 2009; UNODC, 2015a).

ATS are psycho-stimulants that are relatively easy to synthesize and increasingly injected in settings previously dominated by opiates (Bao, et al., 2012; Booth, et al., 2008; Grund, et al., 2009). ATS have been associated with greater sexual risk, including multiple sex partners and unprotected sex, which may compound the risks of HIV acquisition among PWID (Baker, et al., 1994; Booth, et al., 2008; Darke, et al., 2008; Gleghorn, et al., 1998; Molitor, et al., 1999; Molitor, et al., 1998). ATS injection has also been associated with

more frequent injecting, needle/syringe sharing and HIV infection in settings where PWID also injected other drugs (Braine, et al., 2005; Hayashi, et al., 2011; Kozlov, et al., 2006; Tavitian-Exley, et al., 2017) but not when stimulants were reported as main injection drug (Booth, et al., 2008; Kral, et al., 1998; Swe, et al., 2012; Talu, et al., 2010). Few studies have examined drug use patterns by main drug injected and potential associations with risk behaviours and HIV and HCV infection in Eastern European settings (Booth, et al., 2008; Harrell, et al., 2012; Talu, et al., 2010; Tavitian-Exley, et al., 2015) and the relevance of ATS injection in shaping these epidemics remains unclear (EMCDDA, et al., 2011; EMCDDA, et al., 2014).

Our aim is to assess whether primarily injecting ATS as compared to opioids (heroin, synthetic heroin or methadone) is associated with increased injecting and sexual risk behaviours and HIV status among PWID in Kohtla-Järve (Estonia) and St. Petersburg (Russian Federation), two East European locales with significant and epidemiologically similar HIV epidemics.

METHODS

Study population

Integrated biological and behavioural surveys of HIV prevalence were conducted among PWID in Kohtla-Järve between May and July 2012, and in St. Petersburg from November 2012 to June 2013. These surveys used comparable recruitment criteria and respondent-driven sampling (RDS) survey methodology and have been reported on and described previously (Cepeda, et al., 2015; Dukhovlinova, et al., 2015; Heimer, et al., 2015; Tavitian-Exley, et al., 2017; Uusküla, Raag, et al., 2015). Briefly, RDS starts with a diverse sample of seeds (6 seeds in Kohtla-Järve and 12 seeds in different districts of St. Petersburg, subsequently increased to 16 to cover key districts and compensate for unproductive seeds). Seeds were selected through needle/syringe programmes (NSP) to represent a range of demographic and drug profiles; interviews and testing were conducted in fixed (Kohtla-Järve) and mobile clinics (St Petersburg). Each seed and subsequent participants were given an opportunity to recruit up to three PWID until a predetermined sample size was reached. Men and women aged 18 years or over, who had injected drugs in the past 30 days, lived in Kohtla-Järve or St. Petersburg and provided informed consent for the study were eligible. Eligibility was verified by the presence of injection marks and questions on injection practices before the start of the interview.

Measures

Information on demographic and social factors, injection and sexual risk behaviours and access to harm reduction services were recorded by trained fieldworkers in a structured confidential interviewer-administered questionnaire, using standardised study items and questions from established survey instruments (e.g. WHO Drug Injecting study Phase II survey v2b) (Des Jarlais, et al., 2006; Uusküla, Raag, et al., 2015).

HIV, HCV and HSV status

HIV sero-status was assessed using an HIV Antigen/Antibody Combo Assay (ADVIA Centaur, Siemens Healthcare Diagnostics) and HIV I/II Score line assay confirmatory test (INNO LIA, Fujirebio Europe) in Kohtla-Järve; rapid oral HIV testing was conducted in St. Petersburg using OraQuick ADVANCE® Rapid HIV-1/2 Antibody Tests (OraSure Technologies Inc.) and confirmed at the City AIDS Centre (Uusküla, Raag, et al., 2015). In Kohtla-Järve only, HCV and Herpes Simplex Virus (HSV) reactivity were measured using commercially available kits for antibodies to HCV (Murex anti-HCV v 4.0) and HSV-2 (HSV-2 IgG ELISA, IBL International GmbH).

The primary drug injected was categorised into mutually exclusive groups of primary ATS or primary opioid-injectors, based on the survey item *main drug injected in the past 4 weeks*. Injectors reporting no or “other” primary drug were compared with the rest of the sample and examined in descriptive analysis (and excluded in regression modelling).

Demographic and contextual variables

Demographic and contextual variables included age, sex, ethnicity, highest level of education completed (basic education/secondary and above), main source of income, living arrangements (stable/unstable), past month contact with an NSP, past year drug treatment (opioid substitution in Kohtla-Järve; any drug treatment in St. Petersburg) and having needles or syringes confiscated by the police.

Injecting and sexual risk behaviours

The behavioural variables examined included injecting-risk (using a 30 day recall period), sexual-risk behaviours (using a 6 month recall period) and serological markers for HIV (and in Kohtla-Järve only, HCV and HSV). Injecting behaviours of interest were past month injecting frequency (\geq daily injecting vs. $<$ daily injecting), intensity of injection on the last day injected (≥ 2 injections/day vs. <2 injections/day), injecting with used needles/syringes (sharing), sharing drug paraphernalia, back-loading (filling a syringe from a used syringe) and polydrug use (injecting a main drug and at least one other drug

in the last month). Sexual risk behaviours included having a sex partner (regular or casual) who injected drugs, having been paid for sex (i.e. receiving money or drugs for sex ever), multiple sex partners (≥ 2 sex partners in last 6 months) and consistent condom use (i.e. always) with sex partners. The variable “any sex in the last six months” was used to exclude non-sexually active PWID.

Statistical analyses

Descriptive statistics are presented for Kohtla-Järve and St. Petersburg separately (RDS-adjusted estimates, using the RDS-II estimator, are presented in supplementary material) (Volz, et al., 2008; Volz, et al., 2012; White, et al., 2015).

Two sets of logistic regressions were performed. The first set examined the determinants of ATS- and opioid-injection. In the second set, we assessed whether ATS was associated with a) injecting-risk behaviours, b) sexual-risk behaviours, and c) HIV, HCV and HSV prevalence. For both sets of regressions, we generated univariate and multivariable estimates (Kirkwood, et al., 2003; UCLA Statistical Consulting Group, 2015). Results for the multivariable regressions were adjusted for age, sex, education, living arrangements and duration of injecting. In the second set of regressions, we additionally adjusted for contact with NSP and drug treatment. These variables were judged to be important potential confounders, based on published evidence and our conceptual framework (Supplementary material: Figure S.1) (Lemstra, et al., 2012; Marshall, et al., 2008; Marshall, et al., 2011; Poundstone, et al., 2004; Tavitian-Exley, 2016). Variables were examined for collinearity and omitted from the model if their variance inflation factor (VIF) was > 2.5 . A complete case analysis was used and observations with missing values were omitted. Odds ratios (OR) with 95% confidence intervals (95%CI) were calculated while also adjusting for clustering of observations by recruitment seed (Kirkwood, et al., 2003). Clusters were defined by a recruitment chain started by a given seed to account for the possibility that participants may be more likely to recruit other PWID with similar characteristics. This was achieved using the `svy` command in Stata (v.13.1) as in univariate analysis (Hosmer, 2000; Kirkwood, et al., 2003; StataCorp., 2013; UCLA Statistical Consulting Group, 2015). Sampling weights were not taken into account in the regressions as their use is often unwarranted for causal inference (Solon, et al., 2013).

Ethics

Ethical approval was obtained from the Ethics Review Board of the University of Tartu (Estonia), the Institutional Review Board at NGO Stellit in St. Petersburg (Russian Federation), and the Human Investigation Committee at Yale University (USA).

RESULTS

Characteristics of study sample

Our study included 591 PWID in Kohtla-Järve and 811 in St. Petersburg (Table 2; supplementary material Table S.1). In both cities, the majority of PWID were male, Russian-speaking or reporting stable living conditions (Table 2; Table S.1). However, more PWID in Kohtla-Järve were under the age of 30 and salaried or with a regular job than in St. Petersburg.

The primary drug commonly injected by PWID in both cities was a synthetic opioid (i.e. illicitly-manufactured fentanyl congeners in Kohtla-Järve, heroin and methadone produced in illicit laboratories in St. Petersburg) while ATS was the second most common drug class. Opioids were the primary drug for 61% of PWID in Kohtla-Järve and 96% in St. Petersburg; ATS was the main drug for 33% in Kohtla-Järve and 4% in St. Petersburg (6% of PWID in Kohtla-Järve had another or no primary drug).

More PWID had been in contact with a NSP in the past six months in Kohtla-Järve (82%) than in St. Petersburg (16%) and over half had ever received some form of attention for substance use disorder (Table 2). Substitution treatment was reported by 13% of PWID in Kohtla-Järve and detoxification by 11% in St. Petersburg in the past 12 months. Similar proportions of PWID had needles/syringes confiscated by the police in the last six months in both cities, but more PWID reported ever being incarcerated in Kohtla-Järve (55%) than in St. Petersburg (34%).

The majority of PWID in both cities had injected for more than 5 years with a mean age at first injection of just under 19 years (Table S.1). Past-month injection risk behaviours were lower in Kohtla-Järve and up to nine times less frequent than in St. Petersburg. Past-month polydrug use (injection of main and other drug) was equally widespread in both cities (47% and 41%). Fewer PWID in Kohtla-Järve had injected at least daily (24%) than in St. Petersburg (36%), shared needles and syringes (6% and 58%), filled from a used syringe (6% and 53%) or shared drug paraphernalia (7% and 68%).

Over three quarters of PWID in both cities had had sexual intercourse in the last 6 months and over half had a sex partner who injected drugs (55% in Kohtla-Järve and 58% in St. Petersburg). Over a third of PWID had multiple sex partners (34% and 49%) and 7% in Kohtla-Järve and 4% in St. Petersburg respectively had ever been paid for sex. Close to half of PWID always used condoms with casual partners in Kohtla-Järve (58%) and St. Petersburg (48%). Due to the high non-response rate (>50%) for some sexual risk behaviours in Kohtla-Järve, results for this city were not shown, given the high likelihood

of bias; we only analysed the effect of ATS- or opioid-injection for those variables in St. Petersburg (Table S.1). More than half of PWID tested positive for HIV in Kohtla-Järve (61%) and in St. Petersburg (56%). HCV and HSV reactivity measured in Kohtla-Järve only, was 75% and 32%, respectively.

Associations between primary ATS injection and injecting and sexual risk behaviours and serology in Kohtla-Järve

Determinants of ATS injection

Determinants of primary ATS injection are presented in Table 3. In Kohtla-Järve, ATS injectors were younger than opioid injectors with greater odds of being under 30 years of age (adjusted Odds Ratio (aOR) =2.1; 95%CI: 1.4-3.2) and of later injection initiation (aOR: 1.1; 95%CI: 1.0-1.4) than their opioid-injecting peers. The groups did not differ on other demographic characteristics. ATS injectors had lower odds of past-year drug treatment (aOR=0.5; 95%CI: 0.3-0.9), past-month contact with an NSP (aOR=0.2; 95%CI: 0.2-0.6), having needles/syringes confiscated by the police (aOR=0.3; 95%CI: 0.1-0.8) and incarceration (aOR=0.3; 95%CI: 0.2-0.5). They had higher odds of obtaining clean needles/syringes from a pharmacy than an NSP (aOR: 4.9 (95%CI: 3.5-6.9).

Associations with injecting risk behaviours

Primary ATS injection was associated with a number of injecting risk behaviours (Table 4). ATS injectors were more likely to have injected for less than 5 years (aOR=3.5; 95% CI: 1.9-6.2) (Table 4). ATS injection was negatively associated with frequent injecting (aOR=0.3; 95%CI: 0.2-0.6), lifetime needle/syringe-sharing (aOR= 0.3; 95%CI: 0.2-0.5) and back-loading (aOR= 0.4; 95%CI: 0.2-0.8) and was positively associated with polydrug use (aOR=2.0; 95%CI: 1.1-3.5). Past-month unsafe sharing practices such as sharing needles/syringe and drug paraphernalia were generally lower among primary ATS injectors but differences did not reach statistical significance (Table 4).

Associations with sexual risk behaviours

One third of PWID reported multiple sex partners with no significant difference between primary ATS- and opioid-injectors; however ATS injectors had greater odds of ever being paid for sex (aOR=2.6; 95%CI: 1.2-5.7)(Table 5).

Associations with HIV, HCV and HSV prevalence

Primary ATS injectors in Kohtla-Järve had lower odds of testing positive for HIV (aOR=0.6; 95%CI: 0.5-0.8) and having antibodies to HCV (aOR=0.5; 95%CI: 0.3-0.6) in

multivariate analysis, compared to primary opioid injectors (Table 6). HSV antibody status was not associated with ATS injection.

Associations between primary ATS injection and injecting and sexual risk behaviours and HIV in St. Petersburg

Determinants of ATS injection

In St. Petersburg, primary ATS injectors had higher odds of being under 30 years of age (aOR=6.8; 95%CI: 2.8-16.5), female (aOR=1.7; 95%CI: 0.7-4.1), of later initiation to injection (aOR: 1.3; 95%CI: 1.1-1.4) or having unstable living arrangements (aOR=2.2; 95%CI: 1.0-4.6) than opioid injectors (Table 3). Contact with NSP and drug treatment were very low in St. Petersburg and did not differ between ATS- and opioid-injectors. However ATS injectors had greater odds of obtaining needles and syringes from sources other than an NSP (e.g. from friends, other PWID, a drug dealer or in the street) (aOR=35; 95%CI: 2.7-472) and lower odds of having been incarcerated and having needles/syringes confiscated compared to opioid injectors.

Associations with injecting risk behaviours

Primary ATS injectors were more likely to report fewer than 5 years of injecting (aOR=8.3; 95%CI: 2.2-31.6)(Table 4). Several injecting risk behaviours were negatively associated with ATS injection. Primary ATS injectors had lower odds of daily or more frequent injecting (aOR=0.2; 95%CI: 0.1-0.9), injecting more than twice a day (aOR: 0.3; 95%CI: 0.1-0.6), sharing needles and syringes (aOR=0.2; 95%CI: 0.1-0.6), filling a syringe from a used syringe (aOR=0.2; 95%CI: 0.1-0.5) and sharing drug paraphernalia (aOR=0.3; 95%CI: 0.1-0.6) than opioid injectors. Polydrug use was frequent among both ATS- and opioid-injectors but did not differ significantly between the two groups.

Associations with sexual risk behaviours

Almost half of PWID in the Russian city reported multiple sex partners, with no significant difference between ATS- and opioid-injectors (Table 5). In multivariate analysis, ATS injectors had greater odds of being paid for sex (aOR=5.2; 95%CI: 1.0-27.0) and using condoms consistently with casual sex partners (aOR=8.0; 95%CI: 1.1-60.0).

Associations with HIV prevalence

Primary ATS injectors in St. Petersburg had lower odds of testing positive for HIV than PWID injecting opioid (aOR=0.3; 95%CI: 0.1-0.7)(Table 6). HCV and HSV serology were not collected in St. Petersburg.

DISCUSSION

We compared risk behaviours and HIV status among self-identified primary ATS-injectors and opioid-injectors in two settings with severe epidemics of drug use and HIV. Our results suggest that, in both locations, PWID primarily injecting ATS consistently differed on demographic characteristics and reported less or equally risky injecting behaviours compared to those who mainly injected opioids. ATS injectors in both cities were younger but started injecting later than opiate injectors, were more likely to report less than 5 years' injecting and generally at earlier stages in their drug injecting careers. The younger age and later onset of injecting among ATS injectors, suggested the emergence of a different group of PWID who may be at earlier stages of drug dependence and injecting less frequently than their opioid-injecting peers.

In St. Petersburg, contact with NSPs was generally low and did not differ between the two groups; there primary ATS injectors were less likely to have injected with used needles/syringes, shared drug paraphernalia or back-loaded than opiate injectors. However in Kohtla-Järve, where harm reduction services are established, ATS injectors were significantly less likely than opiate injectors to have had contact with NSPs and reported similar prevalence of sharing needles/syringes or drug paraphernalia. They were also less likely to back-load than opiate injectors. That more opiate injectors were in contact with NSPs (and OST) in Kohtla-Järve may have contributed to reducing risky injecting behaviours in this group, thus “levelling” injecting risk between the two drug groups.

Our findings are generally consistent with other studies, where self-reported primary ATS injection was associated with younger age and fewer years of injecting among PWID in Australia (Kaye, et al., 2000), the USA (Braine, et al., 2005), Canada (Fairbairn, et al., 2007), Ukraine and elsewhere in Estonia (Booth, et al., 2008; Talu, et al., 2010) and where primary ATS injectors reported similar or lower frequency of injection (Booth, et al., 2008; Braine, et al., 2005; Darke, et al., 2008; Gleghorn, et al., 1998; Kaye, et al., 2000; Maher, et al., 2007; Talu, et al., 2010) and similar frequency of needle and syringe-sharing than opiate injectors (Gleghorn, et al., 1998; Kaye, et al., 2000; Talu, et al., 2010). That primary ATS injectors in our study reported lower or similar injecting risk

behaviours than opiate injectors contrasts with findings from studies where ATS injection was not defined as main drug and occurred in the presence of other injection drugs, including heroin (Braine, et al., 2005; Crofts, et al., 1997; Hayashi, et al., 2011; Kozlov, et al., 2006; Martin, et al., 2010; Tavitian-Exley, et al., 2017). Injecting risk behaviours and infection risk associated with ATS injection may therefore differ depending on whether PWID inject ATS primarily or inject ATS in addition to or in combination with, addictive opiates such as heroin; these underline the need for consistent definitions and regular monitoring of drugs and drug combinations among PWID.

Both ATS-injectors and opioid-injectors in this study reported similarly high prevalence of sexual risk, including multiple sex partners. However primary ATS injectors were more likely to have ever been paid for sex. ATS may be used to increase energy, stamina, libido and to reduce social and sexual inhibition. Further, injection often occurs with peers or sexual partners, possibly generating more needle-sharing opportunities (Darke, et al., 1995; Klee, 1993). Several other studies of PWID also found positive associations between ATS injection and multiple sex partners, unprotected sex and trading sex for money or drugs (Lorvick, et al., 2006; Molitor, et al., 1999). The frequency of sexual risk in both cities and possible intersection with sex work highlight the potential for sexual transmission of HIV, and an unmet need to engage diverse PWID sub-groups with prevention and risk reduction messages emphasising sexual as well as injecting risks (Lorvick, et al., 2006; Molitor, et al., 1998; Rondinelli, et al., 2009).

Finally, HIV prevalence and HCV in Kohtla-Järve; were lower among ATS- than opioid-injectors, consistent with their shorter, cumulative exposure to risk resulting from younger age and later onset of injecting (Kozlov, et al., 2006; Martin, et al., 2010). These differences in HIV status remained after adjusting for factors such as injecting duration, suggesting that other determinants also play a role. Primary ATS injectors are nevertheless vulnerable to HIV acquisition as a result of their age (and gender in St. Petersburg), high prevalence of sexual risk behaviours and low contact with harm reduction services (i.e. NSP and drug substitution treatment in Kohtla-Järve).

The propensity of ATS injectors in Kohtla-Järve to obtain clean needles and syringes from pharmacies rather than NSPs when the majority of PWID in the sample were in contact with harm reduction services, also suggests they were not being reached (Vorobjov, et al., 2009). Furthermore, the lower odds of substitution treatment among ATS injectors in Kohtla-Järve may be expected since methadone-based substitution treatment has shown to reduce injecting risk behaviours and HIV infection and to support

adherence to highly active anti-retroviral therapy (HAART) among opioid users. However no proven pharmacological treatment exists for ATS-dependent injectors (Ahamad, et al., 2015; MacArthur, et al., 2012; World Health Organization, 2013). Where ATS use is prevalent, services that are tailored to the demographic characteristics and specific needs of ATS injectors and include psychosocial interventions, need to be integrated into harm reduction programmes (Mehrerdi, et al., 2014; UNODC, 2010). The problem of ATS injection in these cities, and globally, requires effective low-threshold services able to engage injectors who are young and/or female, do not fit the drug use “profile” associated with heroin and neither seek nor desire contact with services focused on opioid injectors (Lorvick, et al., 2006; Pates, 2013; Shearer, et al., 2002; Vorobjov, et al., 2009).

Moreover, while few studies have looked at the life course of drug use among primary ATS injectors, especially in resource-limited settings (Brecht, et al., 2008), longitudinal and qualitative research would help to understand how primary ATS-injection and associated behavioural risks evolve over time. Consistent and systematic drug (and polydrug) monitoring would also contribute to better understanding heterogeneity among PWID.

Several limitations need to be acknowledged. First, our findings may only be generalizable to other PWID populations in Estonia or the Russian Federation. Second, obtaining standard probability samples of PWID populations is challenging, due to the hidden nature of this group, their stigmatised behaviours and the absence of a sampling frame. Although RDS surveys have demonstrated the ability to reach hidden population sub-groups, the representativeness of our samples cannot be verified (Abdul-Quader, et al., 2006; Heckathorn, 1997; Johnston, et al., 2010). Third, information on injecting and sexual risk behaviours was collected through self-reports and social desirability bias may affect the results. Self-reporting using interviewer-administered questionnaires has shown reliability in several studies and a 30-day period, as was used in this study, has shown to produce reliable recall on drug use and injecting behaviours among PWID (Darke, 1998; Des Jarlais, et al., 1999; Napper, et al., 2010). Additionally, the small number of ATS injectors in St. Petersburg resulted in wide confidence intervals that limited our analyses for this city. Finally, given the dynamic nature of drug use, high prevalence of polydrug use in this population and shorter recall, it is conceivable that misclassification may have occurred between ATS-and opioid-injectors thus leading to possible bias. However, non-differential misclassification of the exposure generally biased the inferences towards the null and, if this is the case here, our results could be considered conservative (Dosemeci, et al., 1990; Kirkwood, et al., 2003).

The strengths of this study include its large sample size and comparisons of two Eastern European locations reporting a high prevalence of HIV and injection of different drug classes. Recruitment of two large and diverse PWID samples was facilitated by the use of RDS and reported according to the STROBE-RDS statement (Johnston, et al., 2016; White, et al., 2015). We systematically compared PWID injecting different drug classes, using consistent definitions, study methods and tested tools and we highlighted important differences between self-identified primary ATS- and opioid-injectors that are of relevance to policy and programmes.

Primary ATS injectors reported lower or similar injecting risk behaviours, lower HIV prevalence and less engagement with services, than opioid injectors. Both groups had high levels of multiple sex partners but primary ATS injection was associated with paid sex, suggesting overlaps between injecting and sexual risk. Low threshold interventions (e.g. behavioural) and supplies targeting the needs of young stimulant injectors are needed to increase their contact with prevention services and reduce sexual risk behaviours. The coverage of harm reduction services, including sexual risk reduction, needs to be increased significantly in St. Petersburg for all PWID.

Acknowledgments

We are grateful to field staff and peers who conducted the surveys and interviews, to harm reduction programmes ('Me aitamesind' in Kohtla-Järve and 'Diakonia' in St. Petersburg) and to staff who assisted with the identification of seeds and provided facilities for interviewing and serological testing, to NGO "Stellit" for contributing to the survey methodology and to participants for their time.

Author contributions

RH, OL and AU designed the surveys in St. Petersburg and Kohtla-Järve and led surveys and data collection. ITE, MC and MMG developed the research idea and analysis plan, ITE conducted the statistical analyses and drafted the manuscript with MMG, MC and LP. RH, OL and AU provided critical review in the interpretation of results and manuscript. All authors reviewed the manuscript.

Declaration of interest: none declared

Table 1. HIV epidemic, context and response among people who inject drugs in Kohtla-Järve (Estonia) and St. Petersburg (Russia)

Indicator	Kohtla-Järve	St. Petersburg
HIV incidence	22 per 100 person-year (2012) ^(NifHD, 2015; Uusküla, Des Jarlais, et al., 2015) All Estonia: 7.5 per 100 person-year (2011) ^(NifHD, 2015; Uusküla, Des Jarlais, et al., 2015)	14.5 per 100 person-year (2008) ^(Niccolai, et al., 2011) 7.2 per 100 person-year (2010) ^(Kozlov, et al., 2016)
HIV Prevalence	63% (95%CI: 56%-67%) ^(Uusküla, Raag, et al., 2015)	59% (95%CI: 52%-59%) ^(Uusküla, Raag, et al., 2015)
PWID population size estimate	2,000 (range: 700-2,500) All Estonia: 5,362 (range: 3,906–9,837) ^(Uusküla, et al., 2013)	83,120 (95%CI: 77,320 -88,920) ^(Heimer, et al., 2010)
% of population who inject drugs	4.5% (2012) ^(Wu J, et al., under review)	5.5% (2008) ^(Heimer, et al., 2010)
Needle/syringe services (start year)	2004	1996
Needle/syringe services (n, type)	5 outreach, 3 fixed NSP ^(NifHD, 2016)	2 mobile, 2 fixed site services (2015)
Clean syringes per PWID per year	All Estonia: 125 syringes/PWID per year (2011)	n/a
Needle/syringes services provided by	NGOs	City AIDS centre (as of 2015), NGOs
Drug substitution (start year)	2004	OST illegal
Type of drug treatment	Opiate substitution	Detoxification only (21 days)
Coverage (% , n and year)	All Estonia: 15% of PWID (n=919, 2014) ^(NifHD, 2016)	11% PWID registered, % in treatment n/a
Drug treatment services provided by	NGOs, clinics	Centralized, in-patient

Table 1: Reference population for Estonia aged 15-44 years old; for St Petersburg aged 20-45 years old. Data collection years are italicised. HIV= Human Immune deficiency virus. PWID people who inject drugs. CI=Confidence Interval. NSP= Needle and Syringe Programme. OST=Opiate Substitution treatment. NGO= Non-governmental organisation. N/a=not available.

Table 2. Characteristics of sample and by reported primary amphetamine injection in Kohtla-Järve (Estonia) and St. Petersburg (Russian Federation)

CHARACTERISTICS	Kohtla-Järve				St. Petersburg			
	All ⁽¹⁾ PWID Kohtla-Järve	N 591	Primary amphetamine (%) ⁽¹⁾	n 195	All ⁽¹⁾ PWID St. Petersburg	N 811	Primary amphetamine (%) ⁽¹⁾	n 27
DEMOGRAPHIC VARIABLES								
Sex								
Female	26%	155	29.7%	58	22%	180	48.2%	13
Male	74%	434	70.3%	137	78%	631	51.9%	14
Missing		2		0		0		0
Age group								
< 30 years	50%	294	61.5%*	120	30%	241	74.1%*	20
>= 30 years	50%	297	38.5%	75	70%	570	25.9%	7
Missing		0		0		0		0
Ethnicity								
Estonian	12%	66	12.3%	24	0%	0	0%	0
Other	7%	43	5.1%	10	4%	36	3.7%	1
Russian	81%	481	82.6%	161	96%	775	96.3%	26
Missing		1		0		0		0
Education completed								
Basic (9 th grade)/vocational	80%	472	81.0%	158	58%	475	29.6%*	8
Secondary (11 th grade)	19%	116	19.0%	37	30%	243	51.9%	14
Higher (St. Petersburg only)	1%	3	0%	0	12%	93	18.5%	5
Missing		0		0		0		0
Living arrangements								
Unstable (hostel, dormitory, shelter)	40%	238	39.5%	77	36%	292	51.9%*	14
Stable (own or partner's flat/house)	60%	352	60.5%	118	64%	519	48.2%	13
Missing		1		0		0		0
Main income source								
irregular/illicit (SP only)	3%	15	-	-	16%	134	3.7%	1
non-regular/dependant	33%	193	38.5%	74	39%	312	37.0%	10
regular/salaried	64%	375	61.5%	118	45%	362	59.3%	16
Missing		8		3		3		0
CONTEXTUAL VARIABLES								
Lifetime drug treatment								
Ever had drug treatment	55%	324	32.8%*	64	72%	582	25.9%*	7
Never in treatment	45%	267	67.2%	131	28%	229	74.1%	20
Missing		0		0		0		0
Drug/substitution treatment								
Yes (12 months)	13%	75	6.7%*	13	11%	86	14.8%	4
No	87%	516	93.3%	182	89%	724	85.2%	23
Missing		0		0		1		0
NSP programme (4 weeks)								
Contact with NSP	82%	451	66.7%*	124	16%	119	3.7%*	1
No contact with NSP	18%	102	33.3%	62	84%	645	96.3%	26
Missing		38		9		47		0
Source of clean needle/syringes								
Other (friend, dealer, street)	5%	27	66.7%*	124	4%	30	40.7%*	11
Pharmacist/chemist	13%	75	26.3%	49	81%	615	55.6%	15
NSP	82%	451	7.0%	13	16%	119	3.7%	1
Missing		38		9		47		0
Incarceration								
Ever in prison	55%	324	30.3%*	59	34%	274	7.4%*	2
Never in prison	45%	267	69.7%	136	66%	537	92.6%	25
Missing		0		0		0		0
Needles/syringes confiscated								
Had N/S confiscated	31%	404	15.9%*	31	26%	212	0%	0
No N/S confiscated	69%	184	84.1%	164	74%	599	100%	27
Missing		3		0		0		0

Table 2: (1) Column percentage. Crude estimates are presented for Kohtla-Järve and St. Petersburg; adjusted estimates using respondent driven sampling weights (RDS-II, Volz-Heckathorn) are shown in supplementary material as is the number of missing observations. NSP=Needle and syringe programme. HIV= Human Immune deficiency Virus. (2) Drug/substitution treatment in last 12 months refers to opiate drug substitution (OST) in Kohtla-Järve and to detoxification (non-OST) in St Petersburg. *Statistically significant result in comparisons of ATS- and opioid-injectors using Pearson's Chi-squared test for proportions (p-value <0.05) or Fisher's exact when expected cell count is <4.

Table 3. Predictors of primary ATS injection in Kohtla-Järve (Estonia) and St. Petersburg (Russian Federation)

DEMOGRAPHIC CHARACTERISTICS (reference: opioid injectors)	Kohtla-Järve - ATS injectors (n=195) OR ⁽¹⁾ 95% CI	aOR ⁽²⁾ 95%CI	St. Petersburg - ATS injectors (n=27) OR ⁽¹⁾ 95% CI	aOR ⁽²⁾ 95%CI
Sex				
Female	1.3(0.9-1.9)	1.3 (0.9-1.6)	3.4 (1.0-11.5)	1.7 (0.7- 4.1)
Male	<i>ref</i>	<i>ref</i>	<i>ref</i>	<i>ref</i>
Age group				
< 30 years	2.0 (1.3-3.3)	2.1 (1.4-3.2)	7.3 (2.6-20.3)	6.8 (2.8-16.5)
>= 30 years	<i>ref</i>	<i>ref</i>	<i>ref</i>	<i>ref</i>
Ethnicity				
Estonian	1.2 (0.8-1.9)	1.1 (0.6-2.1)	-	-
Other	0.6 (0.2-1.8)	0.6 (0.3-1.1)	0.8 (0.1-9.2)	0.8 (0.1-11.4)
Russian	<i>ref</i>	<i>ref</i>	<i>ref.</i>	<i>ref.</i>
Education completed				
Basic (9 th grade)/vocational	1.1 (0.6-1.8)	1.0 (0.6-1.7)	0.3 (0.05-1.7)	2.6 (1.2-6.7)
Secondary (11 th grade)	<i>ref</i>	<i>ref</i>	1.1 (0.4-3.2)	2.5 (0.7-9.6)
Higher (St. Petersburg only)	-	-	<i>ref</i>	<i>ref</i>
Living arrangements				
Unstable (hostel, dorm, shelter)	0.9 (0.6-1.4)	0.9 (0.6-1.2)	2.0 (0.9-4.1)	2.2 (1.0-4.6)
Stable (own/partner home)	<i>ref</i>	<i>ref</i>	<i>ref</i>	<i>ref</i>
Main income source				
irregular/illicit (SP only)	-	-	0.2 (0.02-1.6)	0.1 (0.05-1.5)
non-regular/dependant	1.3 (0.9 -1.7)	1.0 (0.6 -1.5)	0.7 (0.4-1.5)	0.8 (0.4-1.5)
regular/salaried	<i>ref.</i>	<i>ref.</i>	<i>ref</i>	<i>ref</i>
Age at first injection				
Primary ATS/Opioid	1.1 (1.1-1.2)	1.1 (1.0-1.2)	1.1 (1.0-1.3)	1.3 (1.1-1.4)
CONTEXTUAL VARIABLES				
Lifetime drug treatment				
Ever drug treatment	0.2 (0.2-0.3)	0.3 (0.2-0.5)	0.2 (0.1-0.6)	0.2 (0.1-0.5)
Never in treatment	<i>ref</i>	<i>ref</i>	<i>ref</i>	<i>ref</i>
Drug/substitution treatment				
Yes (12 months)	0.4(0.2-0.7)	0.5 (0.3-0.9)	1.5 (0.4-5.8)	1.9 (0.3-12.7)
No	<i>ref.</i>	<i>ref</i>	<i>ref</i>	<i>ref</i>
NSP programme (4 weeks)				
Contact with NSP	0.2 (0.1-0.3)	0.2 (0.2-0.6)	0.2 (0.05-1.0)	0.4 (0.1-2.2)
No contact with NSP	<i>ref</i>	<i>ref</i>	<i>ref</i>	<i>ref</i>
Source of clean needle/syringes				
Other (friend, dealer, street)	2.4 (0.7-9.9)	2.0 (0.5-7.9)	67 (8.1-544)	35 (2.7-472)
Pharmacist/chemist	6.0 (4.1-8.6)	4.9 (3.5-6.9)	2.9 (0.6-14.2)	1.8 (0.3-9.3)
NSP	<i>ref</i>	<i>ref</i>	<i>ref</i>	<i>ref</i>
Incarceration				
Ever in prison	0.2 (0.1-0.3)	0.3 (0.2-0.5)	0.2 (0.1-0.3)	0.2 (0.1-0.8)
Never in prison	<i>ref</i>	<i>ref</i>	<i>ref</i>	<i>ref</i>
Needles/syringes confiscated				
Had N/S confiscated	0.3 (0.1-0.9)	0.3 (0.1-0.8)	-	<i>Too few obs.</i>
No N/S confiscated	<i>ref</i>	<i>ref</i>		

Table 3. (1) Odds ratio (OR) with 95% confidence intervals (95% CI) refer to primary Amphetamine-Type Stimulant (ATS) versus opioids (reference group is "primary opioid injectors"). (2) Multivariable model for ATS was adjusted for predictors: age, sex, education, duration of injecting (and living arrangements in St. Petersburg). Statistically significant results at the $\alpha < 0.05$ are marked in bold. (3) Needle/syringe programme (NSP), drug treatment and needles/syringes confiscated were adjusted for in models where the outcome was injecting risk. (4) Drug/substitution treatment in last 12 months refers to opiate drug substitution (OST) in Kohtla-Järve and to detoxification (non-OST) in St Petersburg.

Table 4. Association between ATS injection and injecting risk behaviours (Kohtla-Järve, St. Petersburg)

OUTCOMES: INJECTING RISK⁽³⁾ (reference: opioid injectors)	Kohtla-Järve – ATS injectors (n=195)		St. Petersburg – ATS injectors (n=27)	
	OR⁽¹⁾95% CI	aOR⁽²⁾95%CI	OR⁽¹⁾95% CI	aOR⁽²⁾95%CI
Duration of injecting				
≤ 5 years	6.1 (3.8-9.6)	3.5 (1.9-6.2)	14.7 (4.2-51)	8.3 (2.2-31.6)
> 5 years	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Frequency of injecting				
Daily or more	0.2 (0.1-0.5)	0.3 (0.2-0.6)	0.2 (0.1-1.0)	0.2 (0.1-0.9)
Less than daily	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Ever shared needles/syringes				
Yes	0.2 (0.1-0.3)	0.3 (0.2-0.5)	0.05 (0.03-0.1)	0.1 (0.1-0.2)
No	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Injected w/used needles/syringes				
Yes	0.5 (0.1-1.6)	0.5 (0.2-1.5)	0.1 (0.05-0.3)	0.2 (0.1-0.6)
No	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Filled from working syringe (back-loaded)				
Yes	0.4 (0.2-1.2)	0.4 (0.2-0.8)	0.1 (0.1-0.3)	0.2 (0.1-0.5)
No	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Shared drug paraphernalia				
Yes	0.5 (0.2-1.3)	0.5 (0.3-1.2)	0.2 (0.1-0.3)	0.3 (0.1-0.6)
No	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Any polydrug use (any)				
≥ 2 drugs	2.2 (1.5-3.4)	2.0 (1.1-3.5)	0.7 (0.4-1.2)	0.7 (0.4-1.2)
Main drug only	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>

Table 4. (1) Odds ratio (OR) with 95% confidence intervals (95% CI) refer to primary Amphetamine-Type Stimulant (ATS) versus opioids (reference group). (2) Multivariable models were adjusted for age, sex, education, duration of injecting, needle/syringe programme (NSP), drug/substitution treatment, needles/syringes (N/S) confiscated (and living arrangements in St. Petersburg). Statistically significant results at the $\alpha < 0.05$ are marked in bold. (3) Injecting risk in the last 4 weeks.

Table 5. Associations between ATS injection and sexual risk behaviours (Kohtla-Järve, St. Petersburg)

OUTCOMES: SEXUAL RISK⁽³⁾ (reference: opioid injectors)	Kohtla-Järve - ATS injectors (n=195)		St. Petersburg - ATS injectors (n=27)	
	OR⁽¹⁾95% CI	aOR⁽²⁾95%CI	OR⁽¹⁾95% CI	aOR⁽²⁾95%CI
Any sex in 6 months				
Yes	1.8 (1.0-3.3)	1.7 (0.9-3.3)	1.8 (0.8-3.9)	1.6 (0.7- 3.8)
No	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Regular sex partner injects				
Yes	0.7 (0.4-1.1)	0.8 (0.5-1.9)	0.8 (0.2-3.3)	0.8 (0.2- 3.0)
No	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Casual sex partner injects				
Yes	1.1 (0.8-1.5)	1.8 (0.5-7.1)	0.5 (0.1-2.6)	0.4 (0.1-2.2)
No	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Was ever paid for sex				
Yes	1.3 (0.9-1.8)	2.6 (1.2-5.7)	12.2 (3.7-40)	5.2 (1.0-27)
No	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Multiple sex partners				
≥ 2 sex partners	0.8 (0.4-1.6)	0.7 (0.3-1.5)	0.9 (0.5-1.8)	0.7 (0.3-1.6)
one sex partner	<i>Ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Condom with regular partner				
Yes (consistent)	0.9 (0.7-1.4)	1.0 (0.7-1.4)	1.9 (0.4-4.6)	1.5 (0.5-4.5)
No	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>
Condom with casual partner				
Yes (consistent)	1.4 (0.6-3.0)	1.2 (0.6-2.6)	10.4 (1.9-57)	8.0 (1.1-60)
No	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>	<i>ref.</i>

Table 5:(1) Odds ratio (OR) with 95% confidence intervals (95% CI) refer to primary Amphetamine-Type Stimulant (ATS) versus opioids (reference group). (2) Multivariable models were adjusted for age, sex, education, duration of injecting, needle/syringe programme (NSP), drug /substitution treatment, needles/syringes (N/S) confiscated (and living arrangements in St. Petersburg). Statistically significant results at the $\alpha < 0.05$ are marked in bold. (3) Sexual risk in the last 6 months.

Table 6. Associations between ATS injection and serological markers (Kohtla-Järve, St. Petersburg)

SEROLOGICAL MARKERS ⁽³⁾ (reference: opioid injectors)	Kohtla-Järve - ATS injectors (n=195)		St. Petersburg - ATS injectors (n=27)	
	OR⁽¹⁾95% CI	aOR⁽²⁾95%CI	OR⁽¹⁾95% CI	aOR⁽²⁾95%CI
HIV status				
Positive	0.4 (0.3-0.6)	0.6 (0.5-0.8)	0.2 (0.1-0.4)	0.3 (0.1-0.7)
Negative	<i>ref</i>	<i>ref</i>	<i>ref</i>	<i>ref</i>
Hepatitis C				
HCV reactive	0.3 (0.2-0.4)	0.5 (0.3-0.6)	Not collected	Not collected
Non-reactive	<i>ref.</i>	<i>ref</i>	-	-
HSV-2 status				
Positive	1.0 (0.6-1.6)	1.2 (0.7-2.0)	Not collected	Not collected
Negative	<i>ref.</i>	<i>ref</i>	-	-

Table 6: (1) Odds ratio (OR) with 95% confidence intervals (95% CI) refer to primary Amphetamine-Type Stimulant (ATS) versus opioids (reference group). (2) Multivariable models were adjusted for age, sex, education, duration of injecting, needle/syringe programme (NSP), drug /substitution treatment, needles/syringes (N/S) confiscated (and living arrangements in St. Petersburg). Statistically significant results at the $\alpha < 0.05$ are marked in bold. HIV= Human Immune deficiency Virus. HCV=Hepatitis C and HSV-2=Herpes Simplex Virus.

References

- Abdul-Quader, A. S., Heckathorn, D. D., McKnight, C., Bramson, H., Nemeth, C., Sabin, K., Gallagher, K., & Des Jarlais, D. C. (2006). Effectiveness of respondent-driven sampling for recruiting drug users in New York City: findings from a pilot study. *Journal of urban health : bulletin of the New York Academy of Medicine*, 83, 459-476.
- Ahamad, K., Hayashi, K., Nguyen, P., Dobrer, S., Kerr, T., Schütz, C. G., Montaner, J. S., & Wood, E. (2015). Effect of low-threshold methadone maintenance therapy for people who inject drugs on HIV incidence in Vancouver, BC, Canada: an observational cohort study. *The Lancet HIV*.
- Baker, A., Kochan, N., Dixon, J., Wodak, A., & Heather, N. (1994). Drug use and HIV risk-taking behaviour among injecting drug users not currently in treatment in Sydney, Australia. *Drug and Alcohol Dependence*, 34, 155-160.
- Bao, Y. P., Liu, Z. M., Lian, Z., Li, J. H., Zhang, R. M., Zhang, C. B., Hao, W., Wang, X. Y., Zhao, M., Jiang, H. F., Yan, S. Y., Wang, Q. L., Qu, Z., Zhang, H. R., Wu, P., Shi, J., & Lu, L. (2012). Prevalence and correlates of HIV and HCV infection among amphetamine-type stimulant users in 6 provinces in China. *Journal of Acquired Immune Deficiency Syndromes*, 60, 438-446.
- Booth, R. E., Lehman, W. E. K., Kwiatkowski, C. F., Brewster, J. T., Sinitsyna, L., & Dvoryak, S. (2008). Stimulant injectors in Ukraine: The next wave of the epidemic? *AIDS and behavior*, 12, 652-661.
- Braine, N., Des Jarlais, D. C., Goldblatt, C., Zadoretzky, C., & Turner, C. (2005). HIV risk behavior among amphetamine injectors at U.S. syringe exchange programs. *AIDS Education and Prevention*, 17, 515-524.
- Brecht, M.-L., Huang, D., Evans, E., & Hser, Y.-I. (2008). Polydrug use and implications for longitudinal research: Ten-year trajectories for heroin, cocaine, and methamphetamine users. *Drug and Alcohol Dependence*, 96, 193-201.
- Cepeda, J. A., Niccolai, L. M., Lyubimova, A., Kershaw, T., Levina, O., & Heimer, R. (2015). High-risk behaviors after release from incarceration among people who inject drugs in St. Petersburg, Russia. *Drug and Alcohol Dependence*, 147, 196-202.
- Crofts, N., & Aitken, C. K. (1997). Incidence of bloodborne virus infection and risk behaviours in a cohort of injecting drug users in Victoria, 1990-1995. *Medical Journal of Australia*, 167, 17-20.
- Darke, S. (1998). Self-report among injecting drug users: A review. *Drug and Alcohol Dependence*, 51, 253-263.
- Darke, S., Kaye, S., McKetin, R., & Duflou, J. (2008). Major physical and psychological harms of methamphetamine use. *Drug and Alcohol Review*, 27, 253-262.
- Darke, S., Ross, J., Cohen, J., Hando, J., & Hall, W. (1995). Injecting and sexual risk-taking behaviour among regular amphetamine users. *AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV*, 7, 19-26.
- Degenhardt, L., Mathers, B. M., Wirtz, A. L., Wolfe, D., Kamarulzaman, A., Carrieri, M. P., Strathdee, S. A., Malinowska-Sempruch, K., Kazatchkine, M., & Beyrer, C. (2014). What has been achieved in HIV prevention, treatment and care for people who inject drugs, 2010-2012? A review of the six highest burden countries. *International Journal of Drug Policy*, 25, 53-60.
- Des Jarlais, D. C., Paone, D., Milliken, J., Turner, C. F., Miller, H., Gribble, J., Shi, Q., Hagan, H., & Friedman, S. R. (1999). Audio-computer interviewing to measure risk behaviour for HIV among injecting drug users: A quasi-randomised trial. *Lancet*, 353 (9165), 1657-1661.
- Des Jarlais, D. C., Perlis, T. E., Stimson, G. V., Poznyak, V., & Collab, W. P. I. D. I. (2006). Using standardized methods for research on HIV and injecting drug use in

- developing/transitional countries: case study from the WHO Drug Injection Study Phase II. *BMC Public Health*, 6.
- Dosemeci, M., Wacholder, S., & Lubin, J. H. (1990). Does nondifferential misclassification of exposure always bias a true effect toward the null value? *American Journal of Epidemiology*, 132, 746-748.
- Dukhovlinova, E., Masharsky, A., Toussova, O., Verevchkin, S., Solovyeva, T., Meringof, M., Paintsil, E., White, E., Barbour, R., Heimer, R., & Kozlov, A. (2015). Two Independent HIV Epidemics in Saint Petersburg, Russia Revealed by Molecular Epidemiology. *AIDS Research and Human Retroviruses*, 31, 608-614.
- El-Bassel, N., Strathdee, S. A., & Sadr, W. M. E. (2013). HIV and people who use drugs in central Asia: Confronting the perfect storm. *Drug and Alcohol Dependence*, 132, S2-S6.
- EMCDDA. (2010). Problem amphetamine and methamphetamine use in Europe. In *Selected Issues*. Lisbon, Portugal: European Monitoring Centre for Drugs and Drug Addiction,.
- EMCDDA. Harm Reduction Overview for Estonia. Retrieved 2016 2016 from www.emcdda.europa.eu/country-data/harm-reduction/estonia.
- EMCDDA, & Europol. (2011). Amphetamines: A European Union perspective in the global context. In J. Publications (Ed.). Lisbon, Portugal: European Monitoring Centre for Drugs and Drug Addiction.
- EMCDDA, & Europol. (2012). Amphetamine: a European Union perspective in the global context. *Trends in Organized Crime*, 15, 222-240.
- EMCDDA, & Škařupová, K. (2014). The levels of use of opioids, amphetamines and cocaine and associated levels of harm: summary of scientific evidence. In J. V. EMCDDA (D. Thanki (Ed.)). Lisbon, Portugal: European Monitoring Centre for Drugs and Drug Addiction,.
- Eritsyanyan, K., Heimer, R., Barbour, R., Odinkova, V., White, E., Rusakova, M. M., Smolskaya, T. T., & Levina, O. S. (2013). Individual-level, network-level and city-level factors associated with HIV prevalence among people who inject drugs in eight Russian cities: a cross-sectional study. *BMJ Open*, 3.
- Estonia Ministry of Health. (2014). HIV in Estonia: Narrative report for Global AIDS Reporting 2014. In National Institute for Health Development (Ed.). Tallinn.
- Fairbairn, N., Kerr, T., Buxton, J. A., Li, K., Montaner, J. S., & Wood, E. (2007). Increasing use and associated harms of crystal methamphetamine injection in a Canadian setting. *Drug and Alcohol Dependence*, 88, 313-316.
- Gleghorn, A. A., Marx, R., Vittinghoff, E., & Katz, M. H. (1998). Association between drug use patterns and HIV risks among homeless, runaway, and street youth in Northern California. *Drug and Alcohol Dependence*, 51 (3), 219-227.
- Grund, J.-P., Zabransky, T., Irwin, K., & Heimer, R. (2009). Stimulant Use in Central and Eastern Europe. In Pates, R. (Richard) and Diane Riley (eds.), *Interventions for Amphetamine Misuse*, Wiley-Blackwell.
- Harrell, P. T., Mancha, B. E., Petras, H., Trenz, R. C., & Latimer, W. W. (2012). Latent classes of heroin and cocaine users predict unique HIV/HCV risk factors. *Drug and Alcohol Dependence*, 122, 220-227.
- Hayashi, K., Wood, E., Suwannawong, P., Kaplan, K., Qi, J., & Kerr, T. (2011). Methamphetamine injection and syringe sharing among a community-recruited sample of injection drug users in Bangkok, Thailand. *Drug and Alcohol Dependence*, 115 (1-2), 145-149.
- Heckathorn, D. D. (1997). Respondent-driven sampling: a new approach to the study of hidden populations. *Social Problems*, 174-199.
- Heimer, R., Lyubimova, A., Barbour, R., & Levina, O. S. (2015). Emergence of methadone as a street drug in St. Petersburg, Russia. *International Journal of Drug Policy*.

- Heimer, R., & White, E. (2010). Estimation of the number of injection drug users in St. Petersburg, Russia. *Drug and Alcohol Dependence*, 109, 79-83.
- Hosmer, D. W. L., S. (2000). *Applied logistic regression*. New York ; Chichester: Wiley.
- Johnston, L. G., Hakim, A. J., Dittrich, S., Burnett, J., Kim, E., & White, R. G. (2016). A Systematic Review of Published Respondent-Driven Sampling Surveys Collecting Behavioral and Biologic Data. *AIDS Behav.*
- Johnston, L. G., & Sabin, K. (2010). Sampling hard-to-reach populations with respondent driven sampling. *Methodological Innovations Online*, 5, 38-48.
- Jolley, E., Rhodes, T., Platt, L., Hope, V., Latypov, A., Donoghoe, M., & Wilson, D. (2012). HIV among people who inject drugs in Central and Eastern Europe and Central Asia: A systematic review with implications for policy. *BMJ Open*, 2.
- Kaye, S., & Darke, S. (2000). A comparison of the harms associated with the injection of heroin and amphetamines. *Drug and Alcohol Dependence*, 58, 189-195.
- Kirkwood, B., & Sterne, J. (2003). Essential Medical Statistics. In (2nd ed ed.). Oxford: John Wiley & Sons.
- Klee, H. (1993). HIV risks for women drug injectors: Heroin and amphetamine users compared. *Addiction*, 88, 1055-1062.
- Kozlov, Shaboltas, A., Toussova, O., Verevokhin, S., Masse, B., Perdue, T., Beauchamp, G., Sheldon, W., Miller, W. C., Heimer, R., Ryder, R. W., & Hoffman, I. F. (2006). HIV incidence and factors associated with HIV acquisition among injection drug users in St Petersburg, Russia. *AIDS*, 20, 901-906.
- Kozlov, Skochilov, R. V., Toussova, O. V., Verevokhin, S. V., Krasnoselskikh, T. V., Malov, S. V., & Shaboltas, A. V. (2016). HIV incidence and behavioral correlates of HIV acquisition in a cohort of injection drug users in St Petersburg, Russia. *Medicine*, 95, e5238.
- Kral, A. H., Bluthenthal, R. N., Booth, R. E., & Watters, J. K. (1998). HIV seroprevalence among street-recruited injection drug and crack cocaine users in 16 US municipalities. *American Journal of Public Health*, 88, 108-113.
- Lemstra, M., Rogers, M., Thompson, A., Moraros, J., & Buckingham, R. (2012). Risk indicators associated with injection drug use in the Aboriginal population. *Aids Care- Psychological and Socio-Medical Aspects of Aids/Hiv*, 24, 1416-1424.
- Lorvick, J., Martinez, A., Gee, L., & Kral, A. H. (2006). Sexual and Injection Risk among Women who Inject Methamphetamine in San Francisco. *Journal of urban health : bulletin of the New York Academy of Medicine*, 83, 497-505.
- MacArthur, G. J., Minozzi, S., Martin, N., Vickerman, P., Deren, S., Bruneau, J., Degenhardt, L., & Hickman, M. (2012). Opiate substitution treatment and HIV transmission in people who inject drugs: Systematic review and meta-analysis. *BMJ (Online)*, 345.
- Maher, L., Li, J., Jalaludin, B., Wand, H., Jayasuriya, R., Dixon, D., & Kaldor, J. M. (2007). Impact of a reduction in heroin availability on patterns of drug use, risk behaviour and incidence of hepatitis C virus infection in injecting drug users in New South Wales, Australia. *Drug and Alcohol Dependence*, 89, 244-250.
- Marshall, B. D. L., Kerr, T., Livingstone, C., Li, K., Montaner, J. S. G., & Wood, E. (2008). High prevalence of HIV infection among homeless and street-involved Aboriginal youth in a Canadian setting. *Harm Reduction Journal*, 5.
- Marshall, B. D. L., Wood, E., Shoveller, J. A., Buxton, J. A., Montaner, J. S. G., & Kerr, T. (2011). Individual, Social, and Environmental Factors Associated with Initiating Methamphetamine Injection: Implications for Drug Use and HIV Prevention Strategies. *Prevention Science*, 12, 173-180.
- Martin, M., Vanichseni, S., Suntharasamai, P., Mock, P. A., van Griensven, F., Pitisuttithum, P., Tappero, J. W., Chiamwongpaet, S., Sangkum, U., Kitayaporn, D., Gurwith, M., & Choopanya, K. (2010). Drug use and the risk of HIV infection amongst injection drug

- users participating in an HIV vaccine trial in Bangkok, 1999-2003. *International Journal of Drug Policy*, 21, 296-301.
- Mathers, B. M., Degenhardt, L., Ali, H., Wiessing, L., Hickman, M., Mattick, R. P., Myers, B., Ambekar, A., & Strathdee, S. A. (2010). HIV prevention, treatment, and care services for people who inject drugs: a systematic review of global, regional, and national coverage. *The Lancet*, 375, 1014-1028.
- Mehrjerdi, Z. A., Abarashi, Z., Noroozi, A., Arshad, L., & Zarghami, M. (2014). Correlates of shared methamphetamine injection among methamphetamine-injecting treatment seekers: the first report from Iran. *International Journal of STD and AIDS*, 25, 420-427.
- Molitor, Ruiz, J. D., Flynn, N., Mikanda, J. N., Sun, R. K., & Anderson, R. (1999). Methamphetamine Use and Sexual and Injection Risk Behaviors Among Out-of-Treatment Injection Drug Users. *The American journal of drug and alcohol abuse*, 25, 475-493.
- Molitor, Truax, S. R., Ruiz, J. D., & Sun, R. K. (1998). Association of methamphetamine use during sex with risky sexual behaviors and HIV infection among non-injection drug users. *Western Journal of Medicine*, 168 (2), 93-97.
- Napper, L. E., Fisher, D. G., Reynolds, G. L., & Johnson, M. E. (2010). HIV Risk Behavior Self-Report Reliability at Different Recall Periods. *AIDS and behavior*, 14, 152-161.
- Niccolai, L. M., Verevchkin, S. V., Toussova, O. V., White, E., Barbour, R., Kozlov, A. P., & Heimer, R. (2011). Estimates of HIV incidence among drug users in St. Petersburg, Russia: continued growth of a rapidly expanding epidemic. *Eur J Public Health*, 21, 613-619.
- NifHD. (2015). HIV in Estonia: Situation, prevention, treatment, and care. In *Narrative report for GARPR 2015* (pp. 23). Tallinn, Estonia: National Institute for Health Development.
- NifHD. (2016). HIV in Estonia: Situation, prevention, treatment, and care. In *Narrative report for Global AIDS Response Progress Reporting* (pp. 29). Tallinn, Estonia: National Institute for Health Development,.
- Pates, R. (2013). Special Issue on injecting, why is this important? *Journal of Substance Use*, 18, 1-2.
- Poundstone, K. E., Strathdee, S. A., & Celentano, D. D. (2004). The social epidemiology of human immunodeficiency virus/acquired immunodeficiency syndrome. *Epidemiologic Reviews*, 26, 22-35.
- Rondinelli, A. J., Ouellet, L. J., Strathdee, S. A., Latka, M. H., Hudson, S. M., Hagan, H., & Garfein, R. S. (2009). Young adult injection drug users in the United States continue to practice HIV risk behaviors. *Drug and Alcohol Dependence*, 104, 167-174.
- Shearer, J., Sherman, J., Wodak, A., & Van Beek, I. (2002). Substitution therapy for amphetamine users. *Drug and Alcohol Review*, 21, 179-185.
- Solon, G., Haider, S., & Wooldridge, J. (2013). What are we weighting for?(NBER Working Paper 18859).
- StataCorp. (2013). Stata survey data reference manual (Release 13). In (Stata: Release 13 ed.). StataCorp LP. College Station, TX: StataCorp.
- Swe, L. A., & Rashid, A. (2012). Prevalence of HIV and the risk behaviours among injecting drug users in Myanmar. *International Journal of Collaborative Research on Internal Medicine and Public Health*, 4 (1), 56-70.
- Talu, A., Rajaleid, K., Abel-Ollo, K., Ruutel, K., Rahu, M., Rhodes, T., Platt, L., Bobrova, N., & Uusküla, A. (2010). HIV infection and risk behaviour of primary fentanyl and amphetamine injectors in Tallinn, Estonia: Implications for intervention. *International Journal of Drug Policy*, 21, 56-63.
- Tavitian-Exley, I. (2016). *The relevance of polydrug use in HIV risk and associated injecting and sexual risk behaviours among people who inject drugs*. Unpublished Doctoral thesis, Imperial College London, United Kingdom.

- Tavitian-Exley, I., Boily, M., Heimer, R., Uusküla, A., Levina, O., & Maheu-Giroux, M. (2017). Polydrug Use and Heterogeneity in HIV Risk Among People Who Inject Drugs in Estonia and Russia: A Latent Class Analysis. *AIDS and behavior*.
- Tavitian-Exley, I., Vickerman, P., Bastos, F. I., & Boily, M.-C. (2015). Influence of different drugs on HIV risk in people who inject: systematic review and meta-analysis. *Addiction*, 110, 572-584.
- UCLA Statistical Consulting Group. Survey Data Analysis in Stata. from http://www.ats.ucla.edu/stat/stata/seminars/svy_stata_8/default.htm.
- UNAIDS. (2016). Global AIDS update. In (2016 ed.). Geneva: United Nations Joint Programme on AIDS.
- UNODC. (2010). Myanmar Situation Assessment on Amphetamine-Type Stimulants. In R. a. T. S. P. UNODC Global Global Synthetics Monitoring: Analyses (Ed.), *Global Global Synthetics Monitoring: Analyses, Reporting and Trends (SMART) Programme*: United Nations Office on Drugs and Crime,.
- UNODC. (2012). World Drug Report 2012. In Research and Trend Analysis Branch (Ed.), *United Nations publication*. Vienna: United Nations Office on Drugs and Crime,.
- UNODC. (2014). World Drug Report 2014. In Research and Trend Analysis Branch (Ed.), *United Nations publication*. Vienna: United Nations Office on Drugs and Crime,.
- UNODC. (2015a). Trends and Patterns of Amphetamine-type Stimulants and New Psychoactive Substances. In UNODC Regional Office for Southeast Asia and the Pacific (Ed.), *The Challenge of Synthetic Drugs in East and South-East Asia and Oceania* (pp. 45). Vienna: United Nations Office on Drugs and Crime,.
- UNODC. (2015b). World Drug Report 2015. In Research and Trend Analysis Branch (Ed.), *United Nations publication*. Vienna: United Nations Office on Drugs and Crime,.
- UNODC. (2016). World Drug Report 2016. In Research and Trend Analysis Branch (Ed.), *United Nations publication*. Vienna: United Nations Office on Drugs and Crime.
- Uusküla, A., Des Jarlais, D. C., Raag, M., Pinkerton, S. D., & Feelemyer, J. (2015). Combined prevention for persons who inject drugs in the HIV epidemic in a transitional country: the case of Tallinn, Estonia. *AIDS Care*, 27, 105-111.
- Uusküla, A., Raag, M., Vorobjov, S., Ruutel, K., Lyubimova, A., Levina, O. S., & Heimer, R. (2015). Non-fatal overdoses and related risk factors among people who inject drugs in St. Petersburg, Russia and Kohtla-Järve, Estonia. *BMC Public Health*, 15.
- Uusküla, A., Rajaleid, K., Talu, A., Abel-Ollo, K., & Des Jarlais, D. C. (2013). A decline in the prevalence of injecting drug users in Estonia, 2005-2009. *International Journal of Drug Policy*, 24, 312-318.
- Volz, E., & Heckathorn, D. D. (2008). Probability Based Estimation Theory for Respondent Driven Sampling. *Journal of Official Statistics*, 24, 79-97.
- Volz, E., Wejnert, C., Cameron, S., Spiller, M., Barash, V., Degani, I., & Heckathorn, D. D. (2012). *Respondent-driven sampling analysis tool (RDSAT) version 7.1*. Ithaca: Cornell University.
- Vorobjov, S., Uusküla, A., Abel-Ollo, K., Talu, A., Ruutel, K., & Des Jarlais, D. C. (2009). Comparison of injecting drug users who obtain syringes from pharmacies and syringe exchange programs in Tallinn, Estonia. *Harm Reduction Journal*, 6.
- Walsh, N., & Maher, L. (2013). HIV and HCV among people who inject drugs in Central Asia. *Drug and Alcohol Dependence*, 132, S37-S40.
- White, R. G., Hakim, A. J., Salganik, M. J., Spiller, M. W., Johnston, L. G., Kerr, L., Kendall, C., Drake, A., Wilson, D., Orroth, K., Egger, M., & Hladik, W. (2015). Strengthening the Reporting of Observational Studies in Epidemiology for respondent-driven sampling studies: "STROBE-RDS" statement. *Journal of Clinical Epidemiology*.
- World Health Organization. (2013). Global update on HIV treatment 2013: results, impact and opportunities.

Wu J, Crawford FW, Raag M, Heimer R, & A., U. (under review). Using data from respondent-driven sampling studies to estimate the number of people who inject drugs: application to the Kohtla-Järve region of Estonia. . *Submitted to PlosOne*.